

RESEARCH ARTICLE



WILEY

Characterization of *Tbr2*-expressing retinal ganglion cells

Ching-Kang Chen¹ | Takae Kiyama² | Nicole Weber¹ | Christopher M. Whitaker² |
Ping Pan² | Tudor C. Badea^{3,4} | Stephen C. Massey^{2,5} | Chai-An Mao^{2,5}

¹Department of Ophthalmology, Baylor College of Medicine, Houston, Texas, USA

²Ruiz Department of Ophthalmology and Visual Science, McGovern Medical School at The University of Texas Health Science Center at Houston (UTHealth), Houston, Texas, USA

³National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA

⁴Research and Development Institute, Transilvania University of Brasov, School of Medicine, Brasov, Romania

⁵The MD Anderson Cancer Center/UTHealth Graduate School of Biomedical Sciences, Houston, Texas, USA

Correspondence

Chai-An Mao, Ruiz Department of Ophthalmology and Visual Science, McGovern Medical School at UTHealth, 6431 Fannin St., MSB 7.024, Houston, TX 77030, USA
Email: chai-an.mao@uth.tmc.edu

Ching-Kang Chen, Department of Ophthalmology, Baylor College of Medicine, Houston, TX 77030, USA
Email: ching-kang.chen@bcm.edu

Funding information

Alice McPherson Retinal Research Foundation; National Eye Institute, Grant/Award Numbers: EY006515, EY021958, EY024376, EY026930, P30EY002520, P30EY028102; Research to Prevent Blindness

Abstract

The mammalian retina contains more than 40 retinal ganglion cell (RGC) subtypes based on their unique morphologies, functions, and molecular profiles. Among them, intrinsically photosensitive RGCs (ipRGCs) are the first specified RGC type emerging from a common retinal progenitor pool during development. Previous work has shown that T-box transcription factor *T-brain 2* (*Tbr2*) is essential for the formation and maintenance of ipRGCs, and that *Tbr2*-expressing RGCs activate *Opn4* expression upon native ipRGC ablation, suggesting that *Tbr2*⁺ RGCs contain a reservoir for ipRGCs. However, the identity of *Tbr2*⁺ RGCs has not been fully vetted. Here, using genetic sparse labeling and single cell recording, we showed that *Tbr2*-expressing retinal neurons include RGCs and a subset of GABAergic displaced amacrine cells (dACs). Most *Tbr2*⁺ RGCs are intrinsically photosensitive and morphologically resemble native ipRGCs with identical retinofugal projections. *Tbr2*⁺ RGCs also include a unique and rare *Pou4f1*-expressing OFF RGC subtype. Using a loss-of-function strategy, we have further demonstrated that *Tbr2* is essential for the survival of these RGCs and dACs, as well as maintaining the expression of *Opn4*. These data set a strong foundation to study how *Tbr2* regulates ipRGC development and survival, as well as the expression of molecular machinery regulating intrinsic photosensitivity.

KEYWORDS

displaced amacrine cells, intrinsically photosensitive retinal ganglion cells, *Opn4*, *Tbr2*

1 | INTRODUCTION

Intrinsically photosensitive retinal ganglion cells (ipRGCs), marked by the expression of *opsin 4* (*Opn4*, aka melanopsin), are the first specified RGC types in developing mouse retina starting from embryonic day 14 (E14) (Mao et al., 2014; McNeill et al., 2011). IpRGCs play essential roles in nonimage-forming vision, such as pupillary light reflex and circadian photoentrainment (Lazzerini Osprei et al., 2017). They also participate in image-forming vision by regulating RGC excitability as well as eye development (Brown et al., 2010; Rao et al., 2013; Schmidt et al., 2014; Sonoda et al., 2018).

Others and we have identified *T-brain 2* (*Tbr2* or *Eomes*) as the key regulator of ipRGC formation (Mao et al., 2008; Mao et al., 2014;

Sweeney et al., 2014). The observation that many *Tbr2*-expressing RGCs activate *Opn4* expression upon genetic ablation of native ipRGCs suggests that *Tbr2*⁺ RGCs contain a reservoir for native ipRGCs (Mao et al., 2014). However, the identity of *Tbr2*⁺ RGCs and the role *Tbr2* plays in these processes remain unanswered. First, an estimated 17.8% of all RGCs were *Tbr2* positive but only ~40% of *Tbr2*⁺ RGCs were thought to be reservoir for ipRGCs (Mao et al., 2014), leaving ~60% of *Tbr2*⁺ RGCs unaccountable. Second, it was shown that all suprachiasmatic nucleus (SCN)-afferent RGCs expressed melanopsin (Baver et al., 2008), inconsistent with the observation that many cells in the *Tbr2*⁺ ipRGC reservoir do not express melanopsin or at levels below the sensitivity of immunohistochemical detection. Third, our previous conclusion that *Tbr2* is

essential for ipRGC survival was drawn from the near-complete loss of melanopsin-positive cells in adult *Opn4^{Cre}:Tbr2^{fl/fl}* retinas. This observation may alternatively be interpreted by the requirement of

Tbr2 for ipRGC development and/or maturation because *Opn4* expression is clearly detectable at early developmental stages. We shall tackle these unanswered questions here.

Figure 1




FIGURE 1 Legend on next page.

To examine the diversity of $Tbr2^+$ retinal cells and $Tbr2$'s roles in ipRGC development, we generated a new knock-in mouse line containing the engineered $Tbr2^{TauGFP-IREScreERT2}$ allele (as shown in Figure 1(a)). Using this line in conjunction with $Rosa^{iAP}$, $Pou4f1^{CKOAP}$, and $Ai9$ reporter lines, we conducted genetic sparse labeling to investigate the diversity of $Tbr2$ -expressing cells. We found that the vast majority of $Tbr2^+$ RGCs are morphologically indistinguishable from the known M1–M6 ipRGCs. Evidently, these $Tbr2^+$ RGCs project to SCN, ventral lateral geniculate nuclei (vLGN), intergeniculate leaflet (IGL), and olivary pretectal nuclei (OPN), which constitute the primary targets of currently known ipRGC retinofugal projections (Delwig et al., 2016; Ecker et al., 2010; Hattar et al., 2006). We also found a small fraction of $Pou4f1$ -expressing $Tbr2^+$ RGCs and identified them morphologically as a unique bushy OFF RGC type with dendritic stratification distinctively different from and below those of the known M1 type. In addition to RGCs which constitute ~46% of $Tbr2^+$ retinal neurons, surprisingly we found that ~54% of $Tbr2^+$ retinal cells contain two types of GABAergic widefield displaced amacrine cells (dACs) and a mediumfield dAC type. These findings explain the mystery in our previous report where only ~40% of $Tbr2^+$ cells were depleted in $Opn4^{Cre}:Tbr2^{ff}$ retinas (Mao et al., 2014). Moreover, by using a loss-of-function approach, we demonstrate that $Tbr2$ is indeed essential for survival of adult $Tbr2^+$ RGCs and dACs. Most importantly, we found that $Opn4$ expression is downregulated prior to the death of $Tbr2$ -deleted RGCs in adult retinas, lending stronger support for the essential role of $Tbr2$ in maintaining $Opn4$ expression.

2 | MATERIALS AND METHODS

2.1 | Animals

$Tbr2^{TauGFP-IREScreERT2}$ mice (described in Figure 1(a)) were generated by inserting a Frt -PGKNeopA- Frt cassette and a synthetic $TauGFP-IREScreERT2$ DNA fragment in the first exon behind the translation initiation codon of $Tbr2$. Gene targeting was conducted as described previously (Mao et al., 2008). Two targeted embryonic stem cell lines were used to generate $Tbr2^{TauGFP-IREScreERT2}$ mice. The $Tbr2^{TauGFP-IREScreERT2}$ mice were subsequently bred to a $R26-FLPeR$ line

to delete the Frt -flanked PGKNeopA cassette to generate the $Tbr2^{TauGFP-IREScreERT2}$ mice. PCR primers $Tbr2f1$ (5'-AAGGGGCACC TACAATCCCGA-3') and $Taur1$ (5'-TTCGTTCCAGGCTTACCATCC-3') were used for genotyping the $Tbr2^{TauGFP-IREScreERT2}$ allele. $Tbr2^{TauGFP-IREScreERT2}$ mice were bred to $Rosa^{iAP}$, $Pou4f1^{CKOAP}$, or $R26-tdTomato$ ($Ai9$) mice to produce double-transgenic mice for sparse labeling, dye filling, and patch clamp recording. Tamoxifen (1–5 consecutive daily injections of 100 μ g/g body weight; Sigma, St. Louis, MO) was injected intraperitoneally for Cre induction. Alternatively, 1 μ l of 4-hydroxytamoxifen (4-OHT; 0.1 mg/ml; Sigma) was injected into the vitreous space of the right eye to activate cells locally. The generation and genotyping of $Rosa^{iAP}$, $Pou4f1^{CKOAP}$, $Ai9$, $R26^{FLPeR}$, $Slc32a1^{Cre}$, and $Opn4^{TauLacZ}$ mice were described previously (Badea, Cahill, et al., 2009; Farley et al., 2000; Hattar et al., 2002; Madisen et al., 2010; Vong et al., 2011). All mice were maintained on C57BL6/129 mixed backgrounds. Mouse lines of either sex at various ages between P21 and 6 months were used. Preweaned animals were housed with their mother; weaned animals were housed in groups of no more than five in individually ventilated cages. All animal procedures followed the US Public Health Service Policy on Humane Care and Use of Laboratory Animals and were approved by the Animal Welfare Committee at The University of Texas Health Science Center at Houston and Institutional Animal Care and Use Committee at Baylor College of Medicine.

2.2 | Immunohistochemical analysis

Retinal sections or flat-mounted retinas were fixed with 4% paraformaldehyde (PFA; Electron Microscopy Sciences, Hatfield, PA), then incubated with the primary antibodies. Alexa Fluor-conjugated secondary antibodies were obtained from Life Technologies, Carlsbad, CA (1:600 dilution). DAPI (2.5 μ g/ml) was used to stain the nuclei. Images were acquired on a Zeiss 780 confocal laser scanning microscope (Carl Zeiss, Thornwood, NY) and exported as TIFF files into Adobe Photoshop (Adobe Systems, San Jose, CA). Cell counting was conducted using the cell counter plugin of NIH ImageJ.

FIGURE 1 Tracing $Tbr2$ -expressing retinal neurons. (a) Genomic structure of $Tbr2$ locus and the engineered allele in the $Tbr2^{TauGFP-IREScreERT2}$ mouse. (b) Double immunofluorescent staining showing $Tbr2$ expression (magenta) in $Tbr2$ -driven GFP^+ cells (green) in an E16.5 $Tbr2^{TauGFP}$ retinal section. (c) Representative immunofluorescent image showing $Tbr2$ -driven GFP^+ cells (green) in a P3 $Tbr2^{TauGFP}$ retinal flatmount. (d) Immunofluorescent image showing $Tbr2$ expression (magenta) in $Tbr2$ -driven GFP^+ cells (green) in a $Tbr2^{TauGFP}$ retinal section. (e) Fluorescent image of a $Tbr2^{TauGFP}$ retinal section showing dense $Tbr2$ -driven GFP signal (green) in ON sublaminae below the cholinergic ChAT bands (blue) in IPL. Note that GFP signal is separated from the ON ChAT band. (f) Representative co-immunofluorescent image showing melanopsin expression (magenta) in $Tbr2$ -driven GFP^+ cells (green) in a P30 $Tbr2^{TauGFP}$ retinal flatmount. (g) Double immunofluorescent staining showing melanopsin expression (magenta) in $Tbr2$ -driven GFP^+ cells (green). (h) Co-immunofluorescent staining on a $Slc32a1^{Cre}:Ai9$ retinal flatmount showing $Tbr2$ expression (blue) with RBPMS⁺ retinal ganglion cells (RGCs) (yellow arrowheads) or tdTomato⁺ RBPMS⁻ displaced amacrine cells (dACs) (white arrowheads). (i) Relative abundance of $Tbr2^+$ RGCs and dACs described in (g). (j) Relative abundance of $Tbr2^+$ RGCs within the entire RGC population (left) and $Tbr2^+$ dACs within the entire dAC population. (k) Co-immunofluorescent staining on a WT retinal flatmount showing melanopsin expression (magenta) with RBPMS⁺ RGCs (green). ChAT, cholinergic acetyltransferase; GCL, ganglion cell layer; INL, inner nuclear layer; IPL, inner plexiform layer. Scale bars: 50 μ m (b–k) [Color figure can be viewed at wileyonlinelibrary.com]

2.3 | Alkaline phosphatase staining

Tamoxifen-treated *Tbr2^{TauGFP/+;Rosa^{iAP/+}}* mice were used for alkaline phosphatase (AP) staining, which was conducted as previously described with minor modifications (Badea et al., 2003; Jamal et al., 2020). Whole eyeballs were fixed with 10% neutrally buffered formalin for 5 min. The retinas were removed and flat mounted on a piece of nitrocellulose membrane, postfixed for 10 min at room temperature, washed twice in phosphate-buffered saline (PBS), and heated in PBS for 30 min in 65°C water bath to inactivate endogenous AP activity. AP staining was performed in 0.1 M Tris (pH 9.5), 0.1 M NaCl, 50 mM MgCl₂, 0.34 g/ml *p*-nitroblue tetrazolium (NBT) chloride, and 0.175 g/ml 5-bromo-4-chloro-3-indolyl phosphate solution for 24–48 h at room temperature with gentle shaking. After staining, retinas were washed three times for 5 min in PBS, postfixed in PBS with 4% PFA briefly, dehydrated through an ethanol series, and then cleared with 2:1 benzyl benzoate/benzyl alcohol.

Montages of the whole retina were acquired on a Zeiss Axio Imager 2 microscope equipped with a motorized xyz drive (Carl Zeiss, White Plains, NY). Individual RGCs were imaged under transmitted light, and z-stacks were collected with a 20× objective (3.1 pixels/μm) at 0.58 μm vertical (z-dimension) intervals. For each RGC, the areas of dendritic arbors and somata, the branching, and the stratification depth within the inner plexiform layer (IPL) were analyzed using Zen lite (Carl Zeiss). Diameters of dendritic arbors and cell bodies were measured for each cell by calculating the diameter of a circle having the same area as a convex polygon, minimally enclosing the dendritic or somatic profile. Individual examples of distinct morphological types for M2, M4, and M5-like cells were traced using Adobe Photoshop or the NeuronJ plugin of ImageJ (Meijering et al., 2004).

2.4 | Tracing RGC axon targets

For inspecting *Tbr2^{CreERT2}*-driven *Rosa^{iAP}* expression in brains, tamoxifen-treated mice were perfused with 4% cold PFA, and then the whole brains were dissected and postfixed with 4% PFA. Cryopreserved brains were sectioned into consecutive 100 μm coronal sections for AP staining.

2.5 | Dye filling

Dye filling was conducted under room light as previously described with minor modifications (Mao et al., 2014). Briefly, retinas were flat-mounted on black membrane filter paper, and perfused with oxygenated Ames' medium (US Biological, Salem, MA) in a recording chamber. The glass microelectrodes were tip-filled with a mixture of 0.5% Lucifer yellow-CH and 3.5% NBT and then backfilled with 3 M LiCl. The tdTomato⁺ RGCs were viewed under an epifluorescence microscope with a 40× water immersion objective and randomly targeted. Cell penetration was verified within the first minute of iontophoresis of Lucifer yellow with negative current (−0.5 nA at 3 Hz). NBT was then injected into cells with a

positive current (+0.5 nA at 3 Hz) for 5–10 min. Tissue was fixed with 4% PFA in 0.1 M PBS for 1 h at room temperature. Filled cells were visualized by incubating with Alexa Fluor 488-conjugated streptavidin (1:1000 dilution) at 4°C overnight. The injected retinas were stained by an anti-GFP antibody to verify the targeted RGCs and an anti-cholinergic acetyltransferase (ChAT) antibody to determine the depth of dendritic stratification of injected cells.

2.6 | Intrinsic membrane property and photosensitivity

Whole cell patch clamp recording was used as described in Kiyama et al. (2019) to test intrinsic photosensitivity of *Tbr2⁺* cells. Retinas from tamoxifen-treated *Tbr2^{CreERT2/+;Ai9}* mice were flat-mounted with ganglion cell side up under dim red light in a perfusion chamber on an Olympus BX51WI fixed-stage microscope (Olympus USA, Central Valley, PA) and supplied (3 ml/min) with a warm (~34°C) carbogenated mammalian Ringer solution containing (in mM) 120 NaCl, 5 KCl, 25 NaHCO₃, 0.8 Na₂HPO₄, 0.1 NaH₂PO₄, 2 CaCl₂, 1 MgSO₄, 10 D-glucose, 0.01 L-AP4, 0.01 AP5, and 0.04 NBQX. Retinal neurons were visualized by differential interference contrast under infrared illumination (> 900 nm) and *Tbr2⁺* cells marked by tdTomato expression were identified by brief 50 ms light pulses (35,000 R*/rod/s) delivered and imaged through the mCherry filter set (Olympus USA). The cumulative total exposure of the retina was less than 1 s to preserve light sensitivity. Once a cell was selected, a 20-min wait time was used to allow dark adaptation before recording. Patch pipettes made from borosilicate glass tubes (8–12 MΩ) were filled with a potassium-based internal solution containing (in mM) 125 K-gluconate, 8 NaCl, 4 ATP-Mg, 0.5 Na-GTP, 5 EGTA, 10 HEPES, and 0.2% biocytin (w/v) with pH adjusted to 7.3 using KOH. The liquid junction potential was 15 mV and not corrected. Current clamp recording was conducted using the AM2400 amplifier (A-M Systems, Sequim, WA, USA) driven by the WinWCP program provided by John Dempster (University of Strathclyde, Glasgow, UK). Signals were filtered at 5 kHz and sampled at 10 kHz. To obtain intrinsic membrane property (IMP), a small negative current was injected immediately after break-in to adjust membrane potential from rest to −64 ± 2 mV. This was followed by five 600-ms serial negative current injections for hyperpolarization and four serial positive current injections for depolarization. The current that hyperpolarizes a cell to 100 ± 5 mV was empirically determined and used to adjust the amplitudes of subsequent serial current steps. Under these conditions, *Tbr2⁺* cells exhibit characteristic IMP profiles that correspond to cells' dendritic morphologies including IPL stratification levels. Intrinsic photoresponse was initiated by a 1-s 470 nm light stimulation (~430 μm in diameter) at intensities of 2.67–6.14 log R*/rod/s. Dendritic morphology of M1-like and other cells were recovered post hoc by staining for biocytin. IPL stratification level was determined in reference to the distance between the two VaChT or ChAT bands with the ON band designated as 0 and the OFF band as −1. Dendritic structure was traced in the Neurolucida 360 and quantified in the Neurolucida Explorer programs (MBF Bioscience, Boston, MA).

2.7 | Antibody characterization

The Pou4f1 antibody (Millipore: Cat# MAB1585; RRID:AB_94166, used at 1:500) is a mouse monoclonal IgG1 antibody made against aa 186–224 of Pou4f1/Brn-3a fused to the T7 gene 10 protein. Immunostaining using this antibody detects RGCs in wild-type mouse retinas but not in Pou4f1-mutant retinas (Xiang et al., 1995). The Tbr2 antibody (Abcam: Cat# ab23345; RRID: AB_778267, used at 1:500) recognizes a synthetic peptide corresponding to mouse Tbr2 aa 650 to the C-terminus. Immunostaining using this antibody detects RGCs in wild-type mouse retina but not in Tbr2-mutant retinas (Mao et al., 2014). The ChAT antibody (Millipore: Cat #AB144p; RRID: AB_2079751, used at 1:400) is a goat polyclonal antibody made against choline acetyltransferase. It has been widely used on immunostaining and western blotting. The GFP antibody (Thermo Fisher, Cat# A10262, RRID:AB_2534023, used at 1:1000) is a chicken IgY polyclonal antibody that recognizes jellyfish *Aequorea Victoria* green fluorescent protein. Its specificity was demonstrated by detection of different targets fused to GFP tag in transiently transfected cells. This antibody is very specific for detecting GFP-tagged proteins. The RBPMS antibody (Phosphosolutions, Cat# 1832-RBPMS, RRID: AB_2492226, used at 1:500) recognizes a synthetic peptide corresponding to aa residues from the N-terminal region of the rat RBPMS. The antibody is prepared from guinea pig serum by affinity purification. It has been widely used by immunofluorescence to detect RGCs (Kwong et al., 2010). The melanopsin antibody (Advanced Targeting Systems, Cat#AB-N38, RRID:AB_1608077, used at 1:1000) recognizes a 15 aa N-terminal peptide of the mouse melanopsin. This antibody is specific for mouse, and is routinely tested by immunofluorescence (Berson et al., 2010).

2.8 | Experimental design and statistical analysis

Mouse lines (*Tbr2^{TauGFP-IRESCreERT2/+}; Rosa^{iAP}*; *Tbr2^{TauGFP-IRESCreERT2/+}; Pou4f1^{CKOAP/+}* and *Tbr2^{TauGFP-IRESCreERT2/+}; Ai9*) of either sex at various ages between P21 to 6 months were used for sparse labeling, dye filling, and patch clamp recording experiments. Cell counting experiments described in Figure 1 were conducted on 3 *Slc32a^{Cre/+}; Ai9* mice. All data are presented as mean \pm SD for each genotype. Morphometric data of Table 1 are presented as mean \pm SEM.

3 | RESULTS

3.1 | Tbr2-driven TauGFP expression recapitulates endogenous Tbr2 expression

In order to trace and sparsely label retinal neurons that express *Tbr2*, we have generated a new knock-in mouse line *Tbr2^{TauGFP-IRESCreERT2}* (labeled as *Tbr2^{TauGFP}* or *Tbr2^{CreERT2}* interchangeably in the text according to its usage relevancy) in which a *TauGFP-IRESCreERT2* dual reporter cassette was inserted in-frame behind the translational start

codon of *Tbr2* (Met, Figure 1(a)). Immunostaining with an anti-GFP antibody detected GFP in E16.5 and P3 developing *Tbr2^{TauGFP/+}* retinas (Figure 1(b,c)). In adult *Tbr2^{TauGFP/+}* retinas, GFP signals were clearly present in Tbr2-expressing cells and the signal weakened when Tbr2 expression levels were barely detectable (white arrowheads in Figure 1(d)). These data validated the usefulness of this mouse line to track endogenous *Tbr2* expression. We then found in adult *Tbr2^{TauGFP}* retinas that GFP signal was conspicuously present at the lower part of IPL below the ON ChAT band (Figure 1(e)). A noticeable though weaker signal was also found in a narrow band lining the border of IPL and inner nuclear layer (INL) (arrowheads in Figure 1(e,g)) similar to that of Opn4/melanopsin staining pattern. By comparing Opn4 and GFP expression on *Tbr2^{TauGFP}* retinal flatmounts and sections, we found that all Opn4-expressing cells express GFP (Figure 1(f,g)) and weak but detectable GFP signal can be seen on melanopsin-marked dendrites (arrowheads in Figure 1(g)), further affirming the utility of this mouse line for studying the relationship between *Tbr2* and melanopsin-expressing ipRGCs.

Previous estimate put Tbr2-expressing retinal cells at approximately 8.9% of all DAPI⁺ nuclei in the ganglion cell layer (GCL) (Mao et al., 2014). The previous conclusion that Tbr2⁺ RGCs must account for 17.8% of all RGCs was based on two assumptions: (1) All Tbr2⁺ cells are RGCs and (2) RGCs account for ~50% of all GCL cells. To test these assumptions, we co-stained GFP and a pan-RGC marker RBPMS (Kwong et al., 2010). Surprisingly, we found that less than half of the GFP⁺ cells expressed RBPMS (data not shown), indicating that only a fraction of Tbr2⁺ cells are RGCs and that the Tbr2⁺RBPMS⁻ cells may be dACs. To verify this, we examined Tbr2 signal with tdTomato-labeled inhibitory neurons in a *Slc32a1^{Cre}; Ai9* mouse (Vong et al., 2011), and found that ~54% of the Tbr2⁺ cells in the GCL are Ai9⁺RBPMS⁻ dACs (53.6 \pm 1.5%, white arrowheads in Figure 1(h,i)), and that the remaining ~46% Tbr2⁺ cells are RBPMS⁺ RGCs (46.4 \pm 1.5%, yellow arrowheads in Figure 1(h,i)). Using RBPMS and *Slc32a1^{Cre}; Ai9* as pan-RGC and dAC readouts, respectively, we conclude that Tbr2⁺ RGCs account for 11.6 \pm 1.0% of all RGCs, and that Tbr2⁺ dACs account for 18.1 \pm 1.3% of all dACs (Figure 1(j)). Moreover, co-immunostaining of melanopsin and RBPMS revealed that ~6.3% (6.26 \pm 1.23%) of RBPMS⁺ RGCs express melanopsin (Figure 1(k)), suggesting that ~50% of Tbr2⁺ RGCs express detectable level of Opn4 under our experimental conditions.

3.2 | Characterization of Tbr2-expressing retinal neurons

To determine the identities of Tbr2-expressing retinal neurons, we bred *Tbr2^{CreERT2}* with the *Rosa^{iAP}* mouse line to produce *Tbr2^{CreERT2/+}; Rosa^{iAP/+}* mice, then injected tamoxifen through the intraperitoneal (IP) route for genetic sparse AP labeling in retinal flatmounts (Figure 2 (a)) (Badea, Hua, et al., 2009; Badea & Nathans, 2004; Jamal et al., 2020). In agreement with the above finding that Tbr2 is expressed in ~18% of dACs, AP-marked retinal neurons were primarily comprised of widefield dACs (dendritic field diameter > 500 μ m)

TABLE 1 Morphometric data and input resistance of Tbr2⁺ RGCs

	Primary dendrites	Branching number	Dendritic area (mm ²)	Dendritic length (mm)	Branch order	Segment size (μm)	Input resistance (MΩ)
M1n (8 ^a)	3.6 ± 0.5	21.3 ± 6.2	0.14 ± 0.02	2.31 ± 0.34	7.9 ± 1.1	32.8 ± 2.8	1239 ± 70 (26 ^b)
M1r (5)	3.2 ± 0.4	18.4 ± 3.7	0.15 ± 0.03	2.39 ± 0.30	7.6 ± 1.4	33.0 ± 2.2	1439 ± 91 (16)
M1s (9)	3.4 ± 0.4	17.2 ± 2.5	0.12 ± 0.02	2.07 ± 0.25	8.0 ± 1.1	37.4 ± 4.8	604 ± 75 (16)
M2-like (3)	5 ± 0.6	32 ± 9.5	0.19 ± 0.05	3.34 ± 0.46	8.3 ± 1.4	27.9 ± 6.5	317 ± 52 (7)
Bistratified (7)	3.6 ± 0.2	37 ± 12.1	0.16 ± 0.05	2.72 ± 0.61	10 ± 2.4	34.7 ± 8.1	473 ± 83 (12)
M4-like (4)	4.5 ± 0.3	31 ± 3.5	0.26 ± 0.07	4.01 ± 0.69	10.8 ± 1.7	34.5 ± 7.5	170 ± 25 (13)
M5-like (9)	3.5 ± 0.4	52 ± 13.9	0.08 ± 0.01	2.22 ± 0.28	12.7 ± 1.8	24 ± 4.8	376 ± 28 (26)

Abbreviations: IMP, intrinsic membrane property; RGC, retinal ganglion cell.

^aThe number of cells traced and analyzed.

^bThe number of cells analyzed including those identified by IMP and dendritic morphology.

among diverse types of RGCs with dendrites entangled within a dense and narrow band in IPL (Figure 2(b) and arrowheads in Figure 2(b')). By breeding *Tbr2*^{CreERT2} into the Ai9 background (Figure 2(c)), we found that tdTomato⁺ cells from tamoxifen-treated *Tbr2*^{CreERT2};Ai9 retinas co-labeled with Tbr2 and Tbr2-driven GFP expression (Figure 2(d,e)). Using neurobiotin (NBT) dye filling to label the tdTomato⁺ cells in tamoxifen-treated *Tbr2*^{CreERT2};Ai9 retinas, we found that these dACs have long dendritic processes that ramify at the ON laminae of the IPL below the ON-ChAT band (Figure 2(f,f') and Figure 3 insets). This stratification pattern is also consistent with the location of GFP signals in cross-sections of *Tbr2*^{TauGFP/+} retinas shown above (Figure 1(d–g)).

Whole-cell current clamp recording of 230 Tbr2⁺ neurons from 36 adult *Tbr2*^{CreERT2};Ai9 retinas found 156 dACs and among them three distinguishable IMP profiles. The most abundant type (65%) has a very high input resistance (1094 ± 38 MΩ, *n* = 75) and asymmetric dendritic morphology similar to the one identified by NBT dye filling (Figure 2(f)), with dendrites stratifying to IPL level of 0.75 ± 0.04 (*n* = 6, Figure 3(a,b) and inset, in reference to designated OFF-ChAT level of –1 and ON-ChAT at 0). The second dAC type (31%) has a much lower input resistance (300 ± 18 MΩ, *n* = 52) but a larger dendritic area and density whose dendrites stratify to 0.77 ± 0.07 IPL level (*n* = 6, Figure 3(c,d) and inset). These two dAC types exhibited no intrinsic photosensitivity. The third type is rarely encountered and is a mediumfield dAC type (4%) with a medium input resistance (748 ± 87 MΩ, *n* = 6) and a statistically significant closer stratification level to the ON-ChAT band at 0.50 ± 0.12 (*n* = 6, *p* < .0001, Figure 3(e,f) and inset). One cell out of six recorded in this mediumfield dAC type exhibited light sensitivity under glutamatergic synaptic blockade (Figure 3(e')), suggesting that it may be connected through electrical synapse to ipRGCs.

It is worth noting that by delivering 4-OHT directly into the vitreous space (1 μl, 0.1 mg/ml), we could label more RGCs than dACs to reduce the interference from dAC-derived dendritic AP signals while characterizing RGC dendritic morphology. In subsequent studies, we shall focus on describing the Tbr2⁺ RGCs.

3.3 | M1-like Tbr2⁺ RGCs

We have examined 291 cells by AP staining from 37 adult retinas and found 50 “M1-like” cells based on stratification of their terminal dendrites to the boundary between IPL and INL, a signature feature of conventional M1 ipRGCs (Figure 4(a–d)). These cells had medium-sized somata (16.16 ± 3.0 μm in diameter) but displayed large variations in dendritic arbor sizes (see green polygons in Figure 4(a–d); arbor diameter: 264.08 ± 48.75 μm) and degree of dendritic branching (23.69 ± 6.20). The heterogeneity reflects discordant observations of M1-ipRGC sizes in the literature (Berson et al., 2010; Ecker et al., 2010; Emanuel et al., 2017; Lee & Schmidt, 2018; Muller et al., 2010; Schmidt & Kofuji, 2011). Most of these M1-like Tbr2⁺ RGCs have two or three primary dendrites, although one cell with a single primary dendrite (red arrowheads in Figure 4(e,e')) and a smaller soma was also encountered (11.3 vs. 16.16 μm). We also detected several Tbr2⁺ RGCs whose somata were in the INL (red arrowheads in Figure 4(f–h)). These cells have variable dendritic arbor sizes as well but all stratify exclusively to the OFF sublaminae of the IPL. In agreement with our previous finding that all displaced ipRGCs express Tbr2 (Mao et al., 2014), these cells appeared to be displaced M1-ipRGCs.

To determine whether morphological heterogeneity of these M1-like Tbr2⁺ RGCs further reflects the heterogeneity in biophysical properties reported for conventional M1 ipRGCs identified in transgenic lines or by retrograde labeling, we performed targeted patch clamp recording of 56 additional Tbr2⁺ cells under glutamatergic synaptic blockade in 13 tamoxifen-treated *Tbr2*^{CreERT2};Ai9 retinas. We found 33 so-called M1-like cells, all with robust intrinsic photosensitivity (Figure 5). Together with cells identified by dendritic morphology only but without testing for their intrinsic photosensitivity, we have in total analyzed 67 cells and placed them in three subgroups based on their IMP profiles, namely, the presence of rebound spiking following current injection, the level of input resistance, and the appearance of a sag upon hyperpolarization (Table 1 and Figure 5). The first and most abundant subgroup, called M1n, is characterized by the presence of a hyperpolarization sag but no rebound spiking (M1n: *n* = 31, 46%) (Figure 5(a)). The second subgroup has both a




FIGURE 2 Identification of *Tbr2*⁺ retinal ganglion cells (RGCs) and displaced amacrine cells (dACs). (a) The genetic sparse labeling system in the *Tbr2*^{CreERT2};*Rosa*^{iAP} mouse line. (b,b') Representative alkaline phosphatase (AP) staining from a flat-mounted P30 *Tbr2*^{CreERT2};*Rosa*^{iAP} retina showing intense AP⁺ *Tbr2*-expressing retinal neurons (b) and the location where their terminal dendrites stratify in inner plexiform layer (IPL) (b'). (c) The fluorescent labeling system of *Tbr2*⁺ neurons in the *Tbr2*^{CreERT2};*Ai9* mouse line. (d,e) Co-immunofluorescent staining on a *Tbr2*^{CreERT2/+};*Ai9* retinal flatmount showing tdTomato expression (magenta, white arrowheads) with *Tbr2* (green) (d) or *Tbr2*-driven GFP (e). (f,f') Representative images showing the morphology of a *Tbr2*-expressing displaced amacrine cell (yellow arrowhead) and a nearby RGC (white arrowhead) revealed by filled neurobiotin (f) and the location where their terminal dendrites stratify in reference to ChAT bands in IPL (f'). ChAT, cholinergic acetyltransferase; NBT, neurobiotin. Scale bars: 500 μ m (b); 20 μ m (d,e); and 100 μ m (f) [Color figure can be viewed at wileyonlinelibrary.com]

hyperpolarization sag and rebound spiking (M1r; $n = 19$, 28%) (Figure 5(b)). The third subgroup has no hyperpolarization sag and with or without rebound spiking (M1s; $n = 17$, 26%) (Figure 5(c)). By tracing representative cells randomly selected from each subgroups and performed quantitative morphometric analysis, we found no

significant differences among them in primary dendrite numbers, branching numbers, dendritic lengths, dendritic field sizes, branching orders, and segment sizes (Table 1). However, cells in the M1s subgroup have a significantly lower input resistance at 604 ± 75 M Ω ($n = 16$) than those of M1n and M1r at 1239 ± 70 M Ω ($n = 26$,




FIGURE 3 Characterization of $Tbr2^+$ displaced amacrine cells (dACs). Representative intrinsic membrane property (IMP) profiles (a,c,e); intrinsic light sensitivity (a',c',e'); dendritic structure (b,d,f); and inner plexiform layer (IPL) stratification levels (insets) of three $Tbr2^+$ -dACs. Membrane potential changes in response to 600 ms current injections of varying amplitude and polarity are shown at left. (a,b) $Tbr2^+$ -dAC1, not intrinsically photosensitive (a'), is the most frequently encountered type (65%) with a relatively high input resistance (a) and characterized by asymmetric short and long dendrites (b) stratifying to 0.75 ± 0.04 IPL position in reference to designated ON-ChAT position at 0 and OFF-ChAT position at -1 (inset). (c,d) $Tbr2^+$ -dAC2 (31%) has the lowest input resistance (c) and is not intrinsically photosensitive (c') with many radiating long dendrites that span the retina (d) and stratify to 0.77 ± 0.07 IPL position (inset). Note the larger currents required to elicit similar amplitude changes in the $Tbr2^+$ -dAC2. (e,f) $Tbr2^+$ -dAC3 is a rarely encountered (4%) mediumfield dAC with discernible IMP (e) and apparent intrinsic photosensitivity (e') with closer stratification level to ON-ChAT band at 0.50 ± 0.12 (inset). Intrinsic photosensitivity was tested under 1-s light step of 5.15 log $R^*/rod/s$. Scale bar: 200 μm [Color figure can be viewed at wileyonlinelibrary.com]

$p < .00001$) and $1439 \pm 91 M\Omega$ ($n = 16$, $p < .00001$), respectively. Because conventional M1-ipRGCs exhibit a broad range of intrinsic photosensitivity (Milner & Do, 2017); therefore, it will be interesting to determine whether input resistance difference observed here may correlate with intrinsic photosensitivity among the M1-like $Tbr2^+$ cells in future studies.

3.4 | Bistratified $Tbr2^+$ RGCs

We identified 73 AP-labeled bistratified RGCs with diverse morphological characteristics (Figure 6). The first type ($n = 14$) had smaller dendritic arbor sizes and extensive branching patterns (Figure 6(a,b)). Their two dendritic arbors stratified next to the inner (OFF) and outer




FIGURE 4 M1-like $Tbr2^+$ retinal ganglion cells (RGCs). (a–h) Representative bright-field images of alkaline phosphatase (AP)-stained $Tbr2^{CreERT2/+};Rosa^{iAP/+}$ retinal flatmounts. Images were focused on the terminal dendrites located in between INL/IPL boundary (labeled as INL) except in (e) where the focus was on the soma in the GCL. The green polygons represent the terminal dendritic arbor. The red arrowhead in E indicates the single primary dendrite of this RGC. Three representative displaced M1-like RGCs (f–h) with somata located in INL (pointed by red arrowheads) were also shown. GCL, ganglion cell layer; INL, inner nuclear layer; IPL, inner plexiform layer. Scale bars: 100 μ m (a–h) [Color figure can be viewed at wileyonlinelibrary.com]

(ON) margins in the IPL, and the dendritic arbors in the ON sublaminae (red polygons) were larger than that in the OFF sublaminae (green polygons) (Figure 6(a,b')). This type of RGC is similar to the recently described M6 ipRGCs from a $Cdh3$ -GFP mouse line (Quattrochi et al., 2019). The other 59 bistratified $Tbr2^+$ RGCs had relatively larger arbors and less dendritic branching. Twenty-five of these cells had dendrites confined almost entirely to the ON sublaminae (Figure 6(c,d)) with only a few straying ones ramifying up into the OFF sublaminae (black arrowheads in Figure 6(c,d')). Eighteen cells had similar dendritic arbors in ON and OFF sublamina (Figure 6(e,e')), while the remaining 16 cells had dendrites stratified mainly to the OFF-layer (Figure 6(f,f')). These larger bistratified RGCs are similar to the M3 ipRGCs described previously (Schmidt & Kofuji, 2011). We have recorded from a total of 14 bistratified $Tbr2^+$ RGCs by current clamp recording. Dendritic morphology were obtained from eight and analyzed as one bistratified group shown in Table 1. We found varying intrinsic photosensitivity, including six cells (43%) without detectable intrinsic photosensitivity under our brightest light stimulation (Figure 6(g)). The IMP profiles of these

bistratified $Tbr2^+$ RGCs are not uniform. Those with larger OFF layer arbors have no hyperpolarization sag ($n = 4$, example in Figure 6(h)), while those with larger ON layer arbors ($n = 3$) have a unique profile marked by a hyperpolarization sag with or without rebound spiking (Figure 6(i–k)). These bistratified $Tbr2^+$ RGCs therefore are diverse not only in dendritic morphology, but also in IMP, and intrinsic photosensitivity.

3.5 | ON-layer stratified $Tbr2^+$ RGCs

Because the two widefield dAC types described above have long and narrowly stratified and massive dendritic network in the ON lamellae, detailed analysis of $Tbr2^+$ RGCs with dendritic stratification to or near this layer was done on 165 cells whose dendrites were not entrenched with others. Their dendritic characteristics were similar to published ON layer stratified M2, M4, and M5 ipRGC subtypes. Below we separate them by the sizes of their dendritic arbors, somata, branching patterns, and IMPs.




FIGURE 5 Intrinsic membrane property (IMP) and intrinsic photosensitivity of M1-like $Tbr2^+$ retinal ganglion cells (RGCs). (a–c) Representative IMP profiles: M1n (a), M1r (b), M1s (c), intrinsic photosensitivity (a'–c') to 1-s light step of indicated intensity in $\log R^*/rod/s$, and dendritic morphology and stratification patterns (a''–c'') from three M1-like $Tbr2^+$ RGC subtypes. See Table 1 for detailed morphometric analysis which reveals no significant difference between the three except that the M1s subtype has a significantly lower input resistance than the other two. Scale bar: 50 μm [Color figure can be viewed at wileyonlinelibrary.com]

3.5.1 | M4-like $Tbr2^+$ RGCs


M4 ipRGCs have been characterized as having the largest dendritic arbor area and soma diameter among all known ipRGCs (Estevez et al., 2012) and a recent study identified them intersectionally in *Opn4^{Cre};Z/EG* mice using coincidental GFP and SMI32 signals (Sonoda et al., 2019). In 37 AP-stained retinas, we analyzed 15 these M4-like $Tbr2^+$ RGCs with large dendritic arbors and somata (arbor diameter: $327.95 \pm 48.51 \mu m$; soma diameter: $19.28 \pm 3.49 \mu m$; $N = 15$) and relatively loose dendritic arbors (Figure 7(a,b')). They displayed less complex branching (branch points: 29.14 ± 3.46) when compared with other ON-layer stratified $Tbr2^+$ RGCs. We encountered 13 of these M4-like cells by current clamp recording and in contrast to M1-like and bistratified RGCs, all exhibited a characteristic IMP profile as

previously described (Figure 7(c)) (Kiyama et al., 2019). We tested and found two cells with intrinsic photosensitivity under our brightest stimulation (Figure 7(d)). We also randomly traced and analyzed four of these M4-like cells and found them to have the largest dendritic arbors among all RGCs in *Tbr2^{CreERT2};Ai9* retinas (Table 1).

3.5.2 | M5-like $Tbr2^+$ RGCs

The native M5 ipRGC subtype has dense, bushy dendritic trees with smaller dendritic arbors when compared to M2 and M4 types and stratifies below the ON ChAT band in the IPL (Ecker et al., 2010; Stabio et al., 2018). We found 40 AP-stained $Tbr2^+$ RGCs with the M5 morphological characteristics (arbor diameter: $158.32 \pm 24.03 \mu m$;

FIGURE 6 Bistratified $Tbr2^+$ retinal ganglion cells (RGCs). (a–f) representative alkaline phosphatase (AP)-stained images of M6-like (a,b') and various types of M3-like (c–f') $Tbr2^+$ RGCs from $Tbr2^{CreERT2/+};Rosd^{iAP/+}$ retinal flatmounts. Images were focused on the ON (a–f) or OFF (a'–f') IPL sublamina. Red polygons represent the terminal dendritic arbor in the ON-layer, while the green polygons represent those in the OFF-layer. The white arrowheads indicate the sparse terminal dendrites in the ON layer, and the black arrowheads indicate those in the OFF-layer. (g) Representative M3- (upper) and M6-like (lower) $Tbr2^+$ RGC intrinsic photoresponses to 1-s light step of indicated intensity in $\log R^*/rod/s$. (h–k) Representative and diverse intrinsic membrane property (IMP) profiles associated with the bistratified $Tbr2^+$ RGCs. The M6 subtype has a distinguishable IMP profile (k) from the M3 subtypes (h–j). Scale bars: 100 μm (a'–f') [Color figure can be viewed at wileyonlinelibrary.com]



soma diameter: $14.38 \pm 1.86 \mu m$, $N = 35$, Figure 8(a,b)). Notably, their dendrites emanated from three to five primary dendrites (3.75 ± 0.7 , $N = 29$) and stratified slightly below the ON ChAT band (depth in IPL: $85.73 \pm 1.48\%$, $N = 20$), forming more extensive branching than other ipRGCs (49.5 ± 9.58 , $N = 20$). We frequently encountered these M5-like $Tbr2^+$ cells by current clamp recording ($N = 28$). Of 10 cells where intrinsic photosensitivity was tested, 3 (30%) had no detectable intrinsic photosensitivity (Figure 8(c)). The IMP profiles of these M5-like $Tbr2^+$ cells appeared diverse but all contained a unique characteristic delayed return to baseline following positive current injections (Figure 8(d–f), arrows). We randomly traced and analyzed nine such M5-like cells and found them to have the smallest dendritic arbors but the highest number of dendritic branching (Table 1).

Fortuitously, in one animal whose right eye was injected with a low dose of 4-OHT ($1 \mu g/\mu l$), we found a single M5-like RGC in the entire retina (Figure 8(g,g')). AP staining on brain sections revealed its retinofugal projection to the IGL (Figure 8(h,i')); vLGN (Figure 8(k,l',n)); and OPN (Figure 8(m,m',o)). Notably, the size and complexity of its axonal arbors were different in different brain regions. In addition, the axon bifurcated and gave rise to several telodendria along its path

(arrowheads in Figure 8(j',m')). Such projection patterns are similar to those obtained from a single M1-ipRGC, which sends collateral axonal inputs to SCN, IGL, and other brain regions (Fernandez et al., 2016). Whether this is a common feature of all ipRGCs is an interesting topic for future investigations.

3.5.3 | M2-like $Tbr2^+$ RGCs

M2 ipRGCs have an intermediate dendritic arbor size between M4 and M5. We analyzed 24 such “M2-like” AP-stained $Tbr2^+$ RGCs (Figure 9(a–c')). They had relative medium dendritic arbors (arbor diameter: $223.16 \pm 27.82 \mu m$; $N = 21$) and somata (soma diameter: $16.95 \pm 2.22 \mu m$; $N = 22$). Their dendrites emanated from three to five primary dendrites (3.96 ± 0.61 , $N = 23$) and branched more variably than other ipRGCs (40.25 ± 14.03 , $N = 20$). These M2-like cells possessed an IMP profile different from those of M4- and M5-like cells, characterized by the lack of rebound depolarization and immediate return to baseline following current injections (Figure 9(d)). We encountered nine M2-like $Tbr2^+$ cells and of three where intrinsic light response




FIGURE 7 Large M4-like *Tbr2*⁺ retinal ganglion cells (RGCs). (a, b) Representative images of alkaline phosphatase (AP)-stained M4-like *Tbr2*⁺ RGCs from *Tbr2*^{CreERT2/+};*Rosa*^{iAP/+} retinal flatmounts. (a', b') Neuronal tracing of the ministacked images shown in (a, b). (c, d) Stereotypic intrinsic membrane property (IMP) profile (c) and representative intrinsic photoresponse of a M4-like *Tbr2*⁺ RGC to 1-s light step of indicated intensity in log R*/rod/s. (d). Scale bars: 100 μm (a, b') [Color figure can be viewed at wileyonlinelibrary.com]

was examined, two were intrinsically photosensitive (66%) (Figure 9 (e)). Three randomly selected M2-like cells were traced and analyzed with data obtained support the notion that dendritic arbor of these cells lied in between the bigger M4-like and the smaller M5-like cells. IMP profile of these M2-like RGC appears stereotypic (Figure 9(d)).

3.6 | Pou4f1-expressing *Tbr2*⁺ RGCs

A small percentage of *Tbr2*-expressing cells expressed RGC marker *Pou4f1* (Figure 10(a)), which is not expressed, or at a very low level, in ipRGCs (Sweeney et al., 2014), suggesting that these cells are different from aforementioned reservoir ipRGCs. To track them, we bred *Tbr2*^{CreERT2} and *Pou4f1*^{CKO} to produce a *Tbr2*^{CreERT2/+};*Pou4f1*^{CKO/+} mouse line to enable sparse AP staining (Figure 10(b)). In all retinas examined thus far, we found low abundance of these cells with 0 to 2 AP-positive RGCs per retina. The most notable morphological characteristics was the dense and bushy dendrites (Figure 10(c–h')) resulting from numerous branching with short terminal dendrites emanating from multiple primary dendrites (4.15 ± 0.36 , $N = 13$). Their

dendritic arbors and soma sizes were relatively smaller (dendritic arbor diameter: 196.5 ± 26.8 μm; soma diameter: 15.54 ± 1.1 μm; $N = 14$) and their terminal dendrites stratified to the OFF sublaminae (depth in the IPL: $12.8 \pm 1.5\%$, $N = 12$). We classified this RGC type into a unique but rare *Tbr2*⁺*Pou4f1*⁺ RGC subtype because we could only characterize three of them in all *Tbr2*^{CreERT2/+};*Rosa*^{iAP} retinas examined. Despite increasing the sample size in current clamp recording, we have yet to encounter this rare *Tbr2*⁺ RGC type and whether it is intrinsically photosensitive remains to be determined.

3.7 | Axonal projection of *Tbr2*⁺ RGCs in brain

To determine the central projection of *Tbr2*⁺ RGCs, we examined AP staining patterns on brain sections from *Tbr2*^{CreERT2/+};*Rosa*^{iAP/+} mice that received five consecutive tamoxifen doses through the IP route. As expected, intense AP staining was detected in SCN (Figure 11(a)), vLGN, IGL, OPN, and medial pretecal nuclei (Figure 11(b, c)). In addition, a few AP⁺ axonal arbors with relatively diffused signals were found in the dorsal lateral geniculate nuclei and lateral dorsal nucleus (Figure 11(b)), superior colliculus (Figure 11(d–e)), peri-supra-optic nucleus (Figure 11(f)), lateral hypothalamus (Figure 11(g1, g2)), and lateral habenula (Figure 11(h)). These data confirm that *Tbr2*⁺ RGCs are not merely morphologically similar to ipRGCs, they also have very similar, if not identical, retinofugal projections (Delwig et al., 2016; Fernandez et al., 2016; Hattar et al., 2002; Li & Schmidt, 2018).

3.8 | *Tbr2* is essential for the survival of *Tbr2*⁺ RGCs and *Opn4* expression in adult retina

Previous data obtained from *Opn4*^{Cre};*Tbr2*^{fx/fx} mice showed that most *Opn4*-expressing ipRGCs had vanished, leading to the conclusion that *Tbr2* is essential for ipRGC survival. However, since *Opn4* is expressed as early as E14, a caveat lurks that the observed ipRGC loss might result from the requirement of *Tbr2* for ipRGC survival and/or maturation during development. It is therefore necessary to determine whether *Tbr2* is essential for ipRGC survival in adult retinas. To do this, we injected AAV2-Cre into the vitreous space of adult (P30) *Tbr2*^{TauGFP/+} (control) and *Tbr2*^{TauGFP/fx} mice, isolated retinas 12 days postinjection, and examined co-immunostaining patterns of *Tbr2*, GFP, and RBPMS (Figure 12(a, b)). We found that while *Tbr2* expression was efficiently removed by AAV2-Cre in *Tbr2*^{TauGFP/fx} retinas after 12 days (d12), many GFP⁺ *Tbr2*-deficient RGCs remained (double white arrowheads in Figure 12(b)). However, at 21 days postinjection, we found that GFP⁺RBPMS⁺ cells in AAV2-Cre treated *Tbr2*^{TauGFP/fx} retinas decreased by approximately 50% (data not shown). At 38 days postinjection (d38), we could barely find GFP⁺RBPMS⁺ RGCs in *Tbr2*-deleted regions (compared Figure 12(c, d)). Ctrl: 55.2 ± 4.8 ; AAV2-Cre: 10.2 ± 1.5 , $P = 2.2E-06$, per unit area 1.0×10^5 μm²). Together, these data indicate that *Tbr2* is indeed essential for the survival of adult *Tbr2*⁺ reservoir ipRGCs as well as dACs, ascertaining our previous conclusions.




FIGURE 8 Small M5-like *Tbr2*⁺ retinal ganglion cells (RGCs). (a,b) Representative alkaline phosphatase (AP)-stained images of M5-like *Tbr2*⁺ RGCs from *Tbr2*^{CreERT2/+};*Rosa*^{iAP/+} retinal flatmounts. (c) Representative intrinsic photoresponse to 1-s 5.15 log R*/rod/s light step. (d–f) Representative and diverse intrinsic membrane property (IMP) profiles of M5-like *Tbr2*⁺ RGCs. Note the characteristic return delay to the baseline following positive current injections (arrows). (g,g') Retina with a single M5-like *Tbr2*⁺ RGC. (h–p) AP-stained brain sections showing multiple axonal projections of the M5-like RGC shown in (g') in several different regions in the brain. (h'–m') Enlarged images of the bracketed regions in (h) to (m), respectively. IGL, intergeniculate leaflet; INL, inner nuclear layer; OPN, olivary pretectal nuclei; vLGN, ventral lateral geniculate nuclei. Scale bars: 100 μ m (a,b,g') [Color figure can be viewed at wileyonlinelibrary.com]

Because there was a lengthy delay between the deletion of *Tbr2* (12 days postinjection) and the disappearance of *Tbr2*-deficient RGCs (38 days postinjection), we reasoned that if *Opn4* expression requires *Tbr2*, *Opn4* should be downregulated while most of the *Tbr2*-deficient RGCs were still present between d12 and d38. To test this, we injected AAV2-Cre into the vitreous space of P21 *Tbr2*^{fx/+}:*Opn4*^{TauLacZ/+} (control) and *Tbr2*^{fx/fx}:*Opn4*^{TauLacZ/+} retinas, then isolated retinas 14 days postinjection and conducted co-immunostaining for *Tbr2*, RBPMS, and *Opn4*-driven LacZ. We detected fewer LacZ-expressing ipRGCs in the *Tbr2*-deleted region in *Tbr2*^{fx/fx}:*Opn4*^{TauLacZ/+} retinas (compared Figure 12(e,f). Ctrl: 6.3 ± 0.6 ; AAV2-Cre: 3 ± 0.7 , $p = .0003$, per unit area $1.0 \times 10^5 \mu\text{m}^2$). Additionally, most of the remaining LacZ⁺ cells expressed *Tbr2*, suggesting that these cells were

not infected by AAV2-Cre. While the loss of *Opn4* expression did not result in ipRGC death, taken together, these data strongly suggest that *Tbr2* is essential for maintaining *Opn4* expression. Alternatively, the loss of *Opn4* expression is secondary to the eventual death of ipRGCs when *Tbr2* is deleted.

4 | DISCUSSION

Tbr2 is an essential transcription factor for many developmental processes including trophoblast development, mesoderm formation, T cell differentiation, induction of anterior visceral endoderm, and neurogenesis of olfactory and cortical neurons. (Arnold et al., 2008;




FIGURE 9 M2-like $Tbr2^+$ retinal ganglion cells (RGCs). (a–c) Representative alkaline phosphatase (AP)-stained images of 3 M2-like $Tbr2^+$ RGCs from $Tbr2^{CreERT2/+};Rosa^{iAP/+}$ retinal flatmounts. (a'–c') Corresponding tracing of ministacked images shown in (a) to (c). (d) Representative stereotypic intrinsic membrane property (IMP) profile of M2-like $Tbr2^+$ RGCs. Note the quick return to baseline and the absence of regenerative changes following large current injections. (e) Representative intrinsic response to 1-s light step of indicated intensity in $\log R^*/rod/s$. Scale bars: 100 μm (a'–c') [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 10 OFF-layer stratified $Tbr2^+$ $Pou4f1^+$ retinal ganglion cells (RGCs). (a) Fluorescent image of a representative wild-type retinal flatmount showing $Tbr2$ expression (red) with $Pou4f1$ -expressing (green) RGCs (white arrowheads). (b) The genetic sparse labeling system in the $Tbr2^{CreERT2};Pou4f1^{CKO}$ mouse line for bright-field identification of this cell type by AP staining. (c–h') Representative bright-field AP-stained images of the $Tbr2^+$ $Pou4f1^+$ RGCs. Scale bars: 100 μm (c'–h'). AP, alkaline phosphatase [Color figure can be viewed at wileyonlinelibrary.com]

Mizuguchi et al., 2012; Nowotschin et al., 2013; Pearce et al., 2003; Russ et al., 2000; Sessa et al., 2008; Sessa et al., 2010). In retina, we were the first to show that it is a direct downstream target of $Pou4f2$, implicating its role as a regulator for RGC subtype development within the hierarchical transcription network controlling RGC development (Mao et al., 2008; Mu et al., 2008). In follow-up studies, $Tbr2$ was

independently shown to be essential for ipRGC formation (Mao et al., 2014; Sweeney et al., 2014). Using genetic loss-of-function strategies, we further demonstrated that $Tbr2$ is indispensable for ipRGC maintenance and hence proposed that $Opn4$ expression may be activated in $Tbr2^+$ RGCs, allowing them to serve as a reservoir for ipRGCs (Mao et al., 2014). In this study, we developed new mouse




FIGURE 11 Retinofugal projections of $Tbr2^+$ retinal ganglion cells (RGCs) in $Tbr2^{CreERT2/+}; Rosa^{iAP}$ mice. (a–g) Alkaline phosphatase (AP)-stained bright-field images showing projections of $Tbr2^+$ RGCs into SCN (a), IGL, vLGN, dLGN, LD (b); OPN and medial pretectal (MPT) (c); SC (d,e); pSON (f); LDH (g1,g2); and LHb (h). dLGN, dorsal lateral geniculate nuclei; IGL, intergeniculate leaflet; LD, lateral dorsal nucleus; LDH, lateral hypothalamus; LHb, lateral habenula; OPN, olivary pretectal nuclei; pSON, peri-supraoptic nucleus; SC, superior colliculus; SCN, suprachiasmatic nuclei; SO, stratum opticum; u/ISGS, upper/lower stratum griseum superficiale; vLGN, ventral lateral geniculate nuclei

resources to address a number of unanswered questions stemming from this theme.

4.1 | $Tbr2^+$ RGCs are ipRGCs

The first and foremost puzzle that we have solved here is why are there so many $Tbr2^+$ cells but only a small fraction express detectable level of *Opn4*? Our current co-immunostaining and genetically sparse labeling studies reveal that approximately 54% of $Tbr2$ -expressing cells in the GCL are dACs and that the $Tbr2^+$ dACs account for approximately 18% of all dACs in the GCL. Based on their unique IMP profiles and dendritic structures, we have identified two widefield types without intrinsic photosensitivity and one mediumfield type of $Tbr2^+$ dACs with light responsiveness under glutamatergic synaptic

blockade. They are all ON layer-stratified. The finding of two widefield $Tbr2^+$ dACs and a mediumfield type is consistent with one recent scRNA-seq profiling study that revealed several AC subtypes with *Tbr2* expression (Yan et al., 2020). While these dACs were not characterized further, the new $Tbr2^{TauGFP-IRESCreERT2}$ mouse generated for this study should facilitate future studies for their functions. In the RGC population, we found that $Tbr2^+$ RGCs account for approximately 12% of all RGCs, much less than previous estimates. The assumption that all $Tbr2^+$ cells were RGCs must therefore be discarded. Moreover, using melanopsin expression together with a pan-RGC marker (RBPMS), we found that melanopsin-expressing cells account for approximately 6% of all RGCs. This number now approximates the number of GFP^+ RGCs found in $Opn4^{Cre};Z/EG$ retinas in our previous analysis (number of GFP^+ : 2028 at 1-month-old vs. 2870 at 10 months) (Mao et al., 2014). However, it still represents only about half of all $Tbr2^+$ RGCs. Based on our current findings that the majority of $Tbr2^+$ RGCs are anatomically indistinguishable from known ipRGC types and with the same retinofugal projections and many possess intrinsic photosensitivity, we have further strengthened the notion that $Tbr2^+$ RGCs indeed serve as a reservoir for native ipRGCs. The simplest reconciling explanation is that some $Tbr2^+$ RGCs express *Opn4* above detection by conventional immunostaining and those whose levels fall below were not accounted for in previous works. This explanation is supported by the fact that ~82% $Tbr2^+$ RGCs we recorded exhibited intrinsic light sensitivity because whole-cell recording is more sensitive than immunostaining for *Opn4* detection. An alternative and more sophisticated explanation, inspired by the literature, is that the expression of *Opn4* within the entire $Tbr2^+$ RGC population may be dynamically regulated by ambient light level. IpRGC number and *Opn4* level can be modulated by light for the building of a more sensitive and broader photoreceptive system in rats (Hannibal, 2006; Hannibal et al., 2005). It is therefore plausible that the $Tbr2^+$ ipRGC reservoir confirmed in this work allows *Opn4* expression to be dynamically regulated to match the need to adapt to distinct lighting conditions in different seasons and/or geographic locations (Peirson et al., 2009). Our data that *Tbr2* is essential for *Opn4* expression supports this notion. Together, these data strongly bolster our conclusion that $Tbr2^+$ RGCs are bona fide ipRGCs and that it would be of interest in the future to study how *Tbr2* regulates *Opn4* expression.

4.2 | *Tbr2* expression marks all known ipRGC subtypes

Within $Tbr2^+$ RGCs, we identified two types of OFF-layer stratified RGCs, several types of bistratified RGCs, and three types of ON-layer stratified RGCs. The predominant type of the OFF-layer $Tbr2^+$ RGCs resembles previously described M1 ipRGCs, which branch sparsely with monostratified dendritic ramifications to the outermost IPL with variable dendritic field sizes and IMP profiles. This M1 type responds to light through a phototransduction mechanism involving members of the Gq family, Phospholipase C beta 4, and *Trpc6* and *trpc7*




FIGURE 12 Tbr2 is essential for maintaining intrinsically photosensitive retinal ganglion cell (ipRGC) survival and Opn4 expression. (a–d) Fluorescent images of Tbr2, GFP, and RBPMS on *Tbr2^{TauGFP/+}* retinas (a,c) and *Tbr2^{TauGFP/fx}* (b,d) that was treated with AAV2-Cre. Retinas were isolated 12 days (a,b) or 38 days (c,d) postinjection. (e,f) Co-immunofluorescent staining of Tbr2, LacZ, and RBPMS on *Tbr2^{fx/+}*; *Opn4^{TauLacZ}* (e) or *Tbr2^{fx/fx}*; *Opn4^{TauLacZ}* (f) retinas. Retinas were isolated 14 days postinjection. Scale bars: 20 μ m [Color figure can be viewed at wileyonlinelibrary.com]

channels (Emanuel et al., 2017; Xue et al., 2011). We have classified the *Tbr2⁺* M1-like ipRGCs here into three arbitrary subtypes based on their IMP profiles: M1n, M1r, and M1s. Although quantitative morphometric analysis found no significant difference in dendritic structure, the M1s subtype has a significantly lower input resistance than M1n and M1r. While the exact relationship between IMP profiles, dendritic morphology, and intrinsic photosensitivity was not

determined further here, these data suggest that IMP may be a useful “on the fly” cell identifier during recording and can therefore be used to complement post hoc morphological identification. The second and much rarer type of OFF-layer stratified RGCs was discovered in *Tbr2^{CreERT2}*; *Pou4f1^{CKO}* retinas. The co-expression of Tbr2 and Pou4f1 marks this unique subtype, which has bushy and small dendritic arbors stratifying into the OFF sublaminae slightly below the IPL/INL

boundary. The overall morphology of this RGC subtype is reminiscent of types “d” and “j” *Pou4f1*⁺ RGCs (Badea & Nathans, 2011). We have also encountered a few of this RGC subtype in *Tbr2*^{CreERT2/+}; *Rosa*^{lAP} retinas, and it is likely that more of this type of RGC were buried in the dense dendritic AP signals from dACs. The IMP and intrinsic photosensitivity of this subtype was not determined, but can be aided in the future by generating new *Pou4f1*-driven reporter mouse lines allowing intersectional fluorescent labeling. We encountered various types of bistratified *Tbr2*⁺ RGCs, which matched well with the published M3 and M6 types. Finally, we found three ON-layer stratified *Tbr2*⁺ RGC types that morphologically correlated well with known M2, M4, and M5 ipRGCs. We have noted that among the three types, M2 and M4 types have stereotypic IMP profiles, while those of the M5 type appear more diverse. It is also noteworthy that phototransduction mechanism in ON-layer-stratified *Tbr2*⁺ RGCs is different from that in the M1-type and also known to be diverse as hyperpolarization activated channels and/or leaky potassium channels are involved (Jiang et al., 2018; Sonoda et al., 2018). The stereotypic IMP profiles of the M2 and the M4 types may aid future investigative efforts. Taken together, the data we have collected confirm that *Tbr2*⁺ RGCs are indeed ipRGCs. In agreement with our past and current findings, two recent profiling studies using single-cell RNA sequencing (scRNA-seq) found in P5 and adult mouse retinas that *Tbr2* expression is enriched in diverse RGC clusters (Rheume et al., 2018; Tran et al., 2019). In P5 retinas, *Tbr2* was found in 7 (out of 40) RGC clusters (clusters 5, 6, 25, 26, 33, 37, 40), while *Opn4* expression was expressed in 6 of the *Tbr2*⁺ clusters (higher levels in clusters 6, 25, and 33, medium to low levels in clusters 5, 26, and 37, and absent in 40) (Rheume et al., 2018). In adult retinas, *Tbr2* was detected in 8 (out of 46) RGC clusters (clusters 7, 8, 22, 29, 31, 33, 40, 43), while *Opn4* expression is enriched in 5 of the *Tbr2*⁺ clusters (22, 31, 33, 40, 43, and not found in 7, 8, 29) (Tran et al., 2019). Furthermore, our birth-dating data suggest that the onset of ipRGC differentiation matches that of *Tbr2*⁺ RGCs (data not shown). Additionally, our unpublished RNA-seq dataset comparing E15.5 WT and *Tbr2*-mutant retinas has uncovered several *Tbr2*-dependent genes that are enriched in distinct ipRGC subtype clusters. Together, these data suggest that the terminal fate of ipRGC subtypes may have been segregated as early as E15.5, and thus *Tbr2*-mediated transcription network is involved in ipRGC subtype specification. Future research that compares the molecular profiles within each *Tbr2*⁺ cell type at different developmental stages will help elucidating the complex genetic network controlling the specification and differentiation of each of the ipRGC subtypes.

4.3 | Role of *Tbr2* in regulating *Opn4* expression

A well-known function of T-box transcription factors is to regulate molecular pathways involved in cell migration and cell–cell and cell–environment interactions (Ciruna & Rossant, 2001; Costello et al., 2011; Nowotschin et al., 2013; Pearce et al., 2003; Russ et al., 2000; Shen et al., 2014; Strumpf et al., 2005). For example,

Tbr1, a paralogue of *Tbr2*, is necessary for neuronal migration, axon projection, establishment of regional and laminar identity in developing cortex and olfactory bulb, as well as RGC dendritic morphogenesis (Hevner et al., 2001; Huang et al., 2014; Kiyama et al., 2019; Liu et al., 2018). In mouse retinas, *Tbr2* expression appears at ~E14 in a subset of RGCs that settled to the innermost region of GCL, suggesting that it plays a role in early steps of RGC development, such as cell fate determination and/or axon pathfinding. However, how *Tbr2* achieves this and what are its downstream targets remain undetermined. Our study here suggests that deleting *Tbr2* in mature retinas leads first to downregulation of *Opn4* expression in ipRGCs, followed by their death, suggesting that in addition to its role in development, *Tbr2* is also essential for maintaining *Opn4* expression and ipRGC survival in adult. Alternatively but less likely, the loss of *Opn4* expression reflects not only the requirement of *Tbr2* presence but also the health of cells in which *Tbr2* expression becomes undetectable. Nonetheless, we think that *Tbr2* uses similar set of downstream targets to exert its functions during development and in the adult stage. In conjunction with our previous study in which *Tbr2* is shown to be a direct downstream target of *Pou4f2* and *Isl1* (Mao et al., 2008; Wu et al., 2015), it now appears that the *Pou4f2*/*Isl1*-*Tbr2*-*Opn4* regulatory hierarchy is necessary for ipRGC formation and survival. However, how the diverse ipRGC subtypes arise from this seemingly simple gene regulatory hierarchy to produce distinct dendritic morphologies and IPL stratification levels, retinofugal projections, IMPs, and molecular profiles remain poorly understood. Nevertheless, the close association of *Tbr2* with all ipRGC subtypes suggests a pivotal role it plays in this process.

In summary, we have demonstrated that *Tbr2*-expressing RGCs are morphologically indistinguishable from known ipRGCs and that *Tbr2* is essential for maintaining *Opn4* expression and the survival of adult ipRGCs as well as three types of dACs. Future studies should investigate *Tbr2*'s co-regulators and downstream effectors in controlling the formation and maintenance of all ipRGC and dAC subtypes. The generation of the new *Tbr2* reporter mouse line provides a unique opportunity to study in depth several biological questions regarding RGC specification, maturation, and terminal differentiation. One example we have shown here is the projection pattern of a single M5-like *Tbr2*⁺ RGC to multiple brain targets including IGL, vLGN, and OPN. This suggests that a single *Tbr2*⁺ RGC can relay photic information that requires different photoresponse kinetics and threshold sensitivities, underlining the multiple roles of a single ipRGC in nonimage-forming vision. The comprehensive brain targeting pattern of a single RGC has been reported for M1-ipRGCs (Fernandez et al., 2016), but such information is still lacking for other ipRGC subtypes. It is of great interest to delineate retinofugal projections of all *Tbr2*⁺ RGC subtypes at the single cell level to fill the gap in the anatomical architecture and hence functional implications for ipRGCs.

ACKNOWLEDGMENTS

This work was supported by grants from the National Institutes of Health-National Eye Institute to C. A. M. (EY024376), C. K. C. (EY026930), C. M. W. (EY021958), S. C. M. (EY006515), and in part

by National Eye Institute Vision Core Grants P30EY028102 (UTHealth) and P30EY002520 (BCM). The Ophthalmology Department of Baylor College Medicine received an unrestricted grant from Research to Prevent Blindness, Inc. C. K. C. is the Alice McPherson Retinal Research Foundation Chair at BCM. The authors acknowledge Dr Jan Parker-Thornburg and Genetically Engineered Mouse Facility at The University of Texas MD Anderson Cancer Center in making *Tbr2*^{TauGFP-IRES-CreERT2/+} mouse line, Dr King-Wai Yau at Johns Hopkins University for sharing *Opn4*^{TauLacZ} mouse, and Dr Kimberly Mankiewicz at UTHealth for proofreading the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

Ching-Kang Chen, Christopher M. Whitaker, Tudor C. Badea, Stephen C. Massey, and Chai-An Mao: Designed experiments. **Ching-Kang Chen, Christopher M. Whitaker, Takae Kiyama, Nicole Weber, Ping Pan, Stephen C. Massey, and Chai-An Mao:** Executed experiments. **Chai-An Mao:** Designed and generated *Tbr2*^{TauGFP-IRES-CreERT2} mice. **Tudor C. Badea:** Provided protocol, mice and reagents. **Ching-Kang Chen and Chai-An Mao:** Wrote the paper.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1002/cne.25208>.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Tudor C. Badea  <https://orcid.org/0000-0003-3086-6713>

Chai-An Mao  <https://orcid.org/0000-0002-9700-6964>

REFERENCES

- Arnold, S. J., Huang, G. J., Cheung, A. F., Era, T., Nishikawa, S., Bikoff, E. K., ... Groszer, M. (2008). The T-box transcription factor Eomes/*Tbr2* regulates neurogenesis in the cortical subventricular zone. *Genes & Development*, 22(18), 2479–2484. <https://doi.org/10.1101/gad.475408>
- Badea, T. C., Cahill, H., Ecker, J., Hattar, S., & Nathans, J. (2009). Distinct roles of transcription factors *brn3a* and *brn3b* in controlling the development, morphology, and function of retinal ganglion cells. *Neuron*, 61(6), 852–864. <https://doi.org/10.1016/j.neuron.2009.01.020>
- Badea, T. C., Hua, Z. L., Smallwood, P. M., Williams, J., Rotolo, T., Ye, X., & Nathans, J. (2009). New mouse lines for the analysis of neuronal morphology using CreER(T)/loxP-directed sparse labeling. *PLoS One*, 4(11), e7859. <https://doi.org/10.1371/journal.pone.0007859>
- Badea, T. C., & Nathans, J. (2004). Quantitative analysis of neuronal morphologies in the mouse retina visualized by using a genetically directed reporter. *The Journal of Comparative Neurology*, 480(4), 331–351. <https://doi.org/10.1002/cne.20304>
- Badea, T. C., & Nathans, J. (2011). Morphologies of mouse retinal ganglion cells expressing transcription factors *Brn3a*, *Brn3b*, and *Brn3c*: Analysis of wild type and mutant cells using genetically-directed sparse labeling. *Vision Research*, 51(2), 269–279. <https://doi.org/10.1016/j.visres.2010.08.039>
- Badea, T. C., Wang, Y., & Nathans, J. (2003). A noninvasive genetic/pharmacologic strategy for visualizing cell morphology and clonal relationships in the mouse. *The Journal of Neuroscience*, 23(6), 2314–2322.
- Baver, S. B., Pickard, G. E., Sollars, P. J., & Pickard, G. E. (2008). Two types of melanopsin retinal ganglion cell differentially innervate the hypothalamic suprachiasmatic nucleus and the olivary pretectal nucleus. *The European Journal of Neuroscience*, 27(7), 1763–1770. <https://doi.org/10.1111/j.1460-9568.2008.06149.x>
- Berson, D. M., Castrucci, A. M., & Provencio, I. (2010). Morphology and mosaics of melanopsin-expressing retinal ganglion cell types in mice. *The Journal of Comparative Neurology*, 518(13), 2405–2422. <https://doi.org/10.1002/cne.22381>
- Brown, T. M., Gias, C., Hatori, M., Keding, S. R., Semo, M., Coffey, P. J., ... Lucas, R. J. (2010). Melanopsin contributions to irradiance coding in the thalamo-cortical visual system. *PLoS Biology*, 8(12), e1000558. <https://doi.org/10.1371/journal.pbio.1000558>
- Ciruna, B., & Rossant, J. (2001). FGF signaling regulates mesoderm cell fate specification and morphogenetic movement at the primitive streak. *Developmental Cell*, 1(1), 37–49.
- Costello, I., Pimeisl, I. M., Drager, S., Bikoff, E. K., Robertson, E. J., & Arnold, S. J. (2011). The T-box transcription factor Eomesodermin acts upstream of *Mesp1* to specify cardiac mesoderm during mouse gastrulation. *Nature Cell Biology*, 13(9), 1084–1091. <https://doi.org/10.1038/ncb2304>
- Delwig, A., Larsen, D. D., Yasumura, D., Yang, C. F., Shah, N. M., & Copenhagen, D. R. (2016). Retinofugal projections from melanopsin-expressing retinal ganglion cells revealed by intraocular injections of Cre-dependent virus. *PLoS One*, 11(2), e0149501. <https://doi.org/10.1371/journal.pone.0149501>
- Ecker, J. L., Dumitrescu, O. N., Wong, K. Y., Alam, N. M., Chen, S. K., LeGates, T., Renna, J. M., Prusky, G. T., Berson, D. M., & Hattar, S. (2010). Melanopsin-expressing retinal ganglion-cell photoreceptors: Cellular diversity and role in pattern vision. *Neuron*, 67(1), 49–60. <https://doi.org/10.1016/j.neuron.2010.05.023>
- Emanuel, A. J., Kapur, K., & Do, M. T. H. (2017). Biophysical variation within the M1 type of ganglion cell photoreceptor. *Cell Reports*, 21(4), 1048–1062. <https://doi.org/10.1016/j.celrep.2017.09.095>
- Estevez, M. E., Fogerson, P. M., Ilardi, M. C., Borghuis, B. G., Chan, E., Weng, S., Auferkorte, O. N., Demb, J. B., & Berson, D. M. (2012). Form and function of the M4 cell, an intrinsically photosensitive retinal ganglion cell type contributing to geniculocortical vision. *The Journal of Neuroscience*, 32(39), 13608–13620. <https://doi.org/10.1523/JNEUROSCI.1422-12.2012>
- Farley, F. W., Soriano, P., Steffen, L. S., & Dymecki, S. M. (2000). Wide-spread recombinase expression using FLPeR (flipper) mice. *Genesis*, 28(3–4), 106–110.
- Fernandez, D. C., Chang, Y. T., Hattar, S., & Chen, S. K. (2016). Architecture of retinal projections to the central circadian pacemaker. *Proceedings of the National Academy of Sciences of the United States of America*, 113(21), 6047–6052. <https://doi.org/10.1073/pnas.1523629113>
- Hannibal, J. (2006). Regulation of melanopsin expression. *Chronobiology International*, 23(1–2), 159–166. <https://doi.org/10.1080/07420520500464544>
- Hannibal, J., Georg, B., Hindersson, P., & Fahrenkrug, J. (2005). Light and darkness regulate melanopsin in the retinal ganglion cells of the albino Wistar rat. *Journal of Molecular Neuroscience*, 27(2), 147–155. <https://doi.org/10.1385/JMN:27:2:147>
- Hattar, S., Kumar, M., Park, A., Tong, P., Tung, J., Yau, K. W., & Berson, D. M. (2006). Central projections of melanopsin-expressing retinal ganglion cells in the mouse. *The Journal of Comparative Neurology*, 497(3), 326–349. <https://doi.org/10.1002/cne.20970>

- Hattar, S., Liao, H. W., Takao, M., Berson, D. M., & Yau, K. W. (2002). Melanopsin-containing retinal ganglion cells: Architecture, projections, and intrinsic photosensitivity. *Science*, 295(5557), 1065–1070. <https://doi.org/10.1126/science.1069609>
- Hevner, R. F., Shi, L., Justice, N., Hsueh, Y., Sheng, M., Smiga, S., ... Rubenstein, J. L. (2001). Tbr1 regulates differentiation of the preplate and layer 6. *Neuron*, 29(2), 353–366.
- Huang, T. N., Chuang, H. C., Chou, W. H., Chen, C. Y., Wang, H. F., Chou, S. J., & Hsueh, Y. P. (2014). Tbr1 haploinsufficiency impairs amygdalar axonal projections and results in cognitive abnormality. *Nature Neuroscience*, 17(2), 240–247. <https://doi.org/10.1038/nn.3626>
- Jamal, L., Kiyama, T., & Mao, C. A. (2020). Genetically directed sparse labeling system for anatomical studies of retinal ganglion cells. *Methods in Molecular Biology*, 2092, 187–194. https://doi.org/10.1007/978-1-0716-0175-4_13
- Jiang, Z., Yue, W. W. S., Chen, L., Sheng, Y., & Yau, K. W. (2018). Cyclic-nucleotide- and HCN-channel-mediated phototransduction in intrinsically photosensitive retinal ganglion cells. *Cell*, 175(3), 652–664 e612. <https://doi.org/10.1016/j.cell.2018.08.055>
- Kiyama, T., Long, Y., Chen, C. K., Whitaker, C. M., Shay, A., Wu, H., Badea, T. C., Mohsenin, A., Parker-Thornburg, J., Klein, W. H., Mills, S. L., Massey, S. C., & Mao, C. A. (2019). Essential roles of Tbr1 in the formation and maintenance of the orientation-selective J-RGCs and a group of OFF-sustained RGCs in mouse. *Cell Reports*, 27(3), 900–915 e905. <https://doi.org/10.1016/j.celrep.2019.03.077>
- Kwong, J. M., Caprioli, J., & Piri, N. (2010). RNA binding protein with multiple splicing: A new marker for retinal ganglion cells. *Investigative Ophthalmology & Visual Science*, 51(2), 1052–1058. <https://doi.org/10.1167/iovs.09-4098>
- Lazzerini Ospri, L., Prusky, G., & Hattar, S. (2017). Mood, the circadian system, and melanopsin retinal ganglion cells. *Annual Review of Neuroscience*, 40, 539–556. <https://doi.org/10.1146/annurev-neuro-072116-031324>
- Lee, S. K., & Schmidt, T. M. (2018). Morphological identification of melanopsin-expressing retinal ganglion cell subtypes in mice. *Methods in Molecular Biology*, 1753, 275–287. https://doi.org/10.1007/978-1-4939-7720-8_19
- Li, J. Y., & Schmidt, T. M. (2018). Divergent projection patterns of M1 ipRGC subtypes. *The Journal of Comparative Neurology*, 526(13), 2010–2018. <https://doi.org/10.1002/cne.24469>
- Liu, J., Reggiani, J. D. S., Laboulaye, M. A., Pandey, S., Chen, B., Rubenstein, J. L. R., Krishnaswamy, A., & Sanes, J. R. (2018). Tbr1 instructs laminar patterning of retinal ganglion cell dendrites. *Nature Neuroscience*, 21(5), 659–670. <https://doi.org/10.1038/s41593-018-0127-z>
- Madisen, L., Zwingman, T. A., Sunkin, S. M., Oh, S. W., Zariwala, H. A., Gu, H., Ng, L. L., Palmiter, R. D., Hawrylycz, M. J., Jones, A. R., Lein, E. S., & Zeng, H. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neuroscience*, 13(1), 133–140. <https://doi.org/10.1038/nn.2467>
- Mao, C. A., Kiyama, T., Pan, P., Furuta, Y., Hadjantonakis, A. K., & Klein, W. H. (2008). Eomesodermin, a target gene of Pou4f2, is required for retinal ganglion cell and optic nerve development in the mouse. *Development*, 135(2), 271–280. <https://doi.org/10.1242/dev.009688>
- Mao, C. A., Li, H., Zhang, Z., Kiyama, T., Panda, S., Hattar, S., Ribelayga, C. P., Mills, S. L., & Wang, S. W. (2014). T-box transcription regulator Tbr2 is essential for the formation and maintenance of Opn4/melanopsin-expressing intrinsically photosensitive retinal ganglion cells. *The Journal of Neuroscience*, 34(39), 13083–13095. <https://doi.org/10.1523/JNEUROSCI.1027-14.2014>
- McNeill, D. S., Sheely, C. J., Ecker, J. L., Badea, T. C., Morhardt, D., Guido, W., & Hattar, S. (2011). Development of melanopsin-based irradiance detecting circuitry. *Neural Development*, 6, 8. <https://doi.org/10.1186/1749-8104-6-8>
- Meijering, E., Jacob, M., Sarria, J. C., Steiner, P., Hirling, H., & Unser, M. (2004). Design and validation of a tool for neurite tracing and analysis in fluorescence microscopy images. *Cytometry. Part A*, 58(2), 167–176. <https://doi.org/10.1002/cyto.a.20022>
- Milner, E. S., & Do, M. T. H. (2017). A population representation of absolute light intensity in the mammalian retina. *Cell*, 171(4), 865–876 e816. <https://doi.org/10.1016/j.cell.2017.09.005>
- Mizuguchi, R., Naritsuka, H., Mori, K., Mao, C. A., Klein, W. H., & Yoshihara, Y. (2012). Tbr2 deficiency in mitral and tufted cells disrupts excitatory-inhibitory balance of neural circuitry in the mouse olfactory bulb. *The Journal of Neuroscience*, 32(26), 8831–8844. <https://doi.org/10.1523/JNEUROSCI.5746-11.2012>
- Mu, X., Fu, X., Beremand, P. D., Thomas, T. L., & Klein, W. H. (2008). Gene regulation logic in retinal ganglion cell development: Isl1 defines a critical branch distinct from but overlapping with Pou4f2. *Proceedings of the National Academy of Sciences of the United States of America*, 105(19), 6942–6947. <https://doi.org/10.1073/pnas.0802627105>
- Muller, L. P., Do, M. T., Yau, K. W., He, S., & Baldrige, W. H. (2010). Tracer coupling of intrinsically photosensitive retinal ganglion cells to amacrine cells in the mouse retina. *The Journal of Comparative Neurology*, 518(23), 4813–4824. <https://doi.org/10.1002/cne.22490>
- Nowotschin, S., Costello, I., Piliszek, A., Kwon, G. S., Mao, C. A., Klein, W. H., Robertson, E. J., & Hadjantonakis, A. K. (2013). The T-box transcription factor Eomesodermin is essential for AVE induction in the mouse embryo. *Genes & Development*, 27(9), 997–1002. <https://doi.org/10.1101/gad.215152.113>
- Pearce, E. L., Mullen, A. C., Martins, G. A., Krawczyk, C. M., Hutchins, A. S., Zediak, V. P., Banica, M., DiCioccio, C., Gross, D. A., Mao, C. A., Shen, H., Cereb, N., Yang, S. Y., Lindsten, T., Rossant, J., Hunter, C. A., & Reiner, S. L. (2003). Control of effector CD8⁺ T cell function by the transcription factor Eomesodermin. *Science*, 302(5647), 1041–1043. <https://doi.org/10.1126/science.1090148>
- Peirson, S. N., Halford, S., & Foster, R. G. (2009). The evolution of irradiance detection: Melanopsin and the non-visual opsins. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 364(1531), 2849–2865. <https://doi.org/10.1098/rstb.2009.0050>
- Quattrochi, L. E., Stabio, M. E., Kim, I., Iardi, M. C., Michelle Fogerson, P., Leyrer, M. L., & Berson, D. M. (2019). The M6 cell: A small-field bistratified photosensitive retinal ganglion cell. *The Journal of Comparative Neurology*, 527(1), 297–311. <https://doi.org/10.1002/cne.24556>
- Rao, S., Chun, C., Fan, J., Kofron, J. M., Yang, M. B., Hegde, R. S., Ferrara, N., Copenhagen, D. R., & Lang, R. A. (2013). A direct and melanopsin-dependent fetal light response regulates mouse eye development. *Nature*, 494(7436), 243–246. <https://doi.org/10.1038/nature11823>
- Rheume, B. A., Jereen, A., Bolisetty, M., Sajid, M. S., Yang, Y., Renna, K., Sun, L., Robson, P., & Trakhtenberg, E. F. (2018). Single cell transcriptome profiling of retinal ganglion cells identifies cellular subtypes. *Nature Communications*, 9(1), 2759. <https://doi.org/10.1038/s41467-018-05134-3>
- Russ, A. P., Wattler, S., Colledge, W. H., Aparicio, S. A., Carlton, M. B., Pearce, J. J., Barton, S. C., Surani, M. A., Ryan, K., Nehls, M., Wilson, V., & Evans, M. J. (2000). Eomesodermin is required for mouse trophoblast development and mesoderm formation. *Nature*, 404(6773), 95–99. <https://doi.org/10.1038/35003601>
- Schmidt, T. M., Alam, N. M., Chen, S., Kofuji, P., Li, W., Prusky, G. T., & Hattar, S. (2014). A role for melanopsin in alpha retinal ganglion cells and contrast detection. *Neuron*, 82(4), 781–788. <https://doi.org/10.1016/j.neuron.2014.03.022>
- Schmidt, T. M., & Kofuji, P. (2011). Structure and function of bistratified intrinsically photosensitive retinal ganglion cells in the mouse. *The Journal of Comparative Neurology*, 519(8), 1492–1504. <https://doi.org/10.1002/cne.22579>

- Sessa, A., Mao, C. A., Colasante, G., Nini, A., Klein, W. H., & Broccoli, V. (2010). Tbr2-positive intermediate (basal) neuronal progenitors safeguard cerebral cortex expansion by controlling amplification of pallial glutamatergic neurons and attraction of subpallial GABAergic interneurons. *Genes & Development*, *24*(16), 1816–1826. <https://doi.org/10.1101/gad.575410>
- Sessa, A., Mao, C. A., Hadjantonakis, A. K., Klein, W. H., & Broccoli, V. (2008). Tbr2 directs conversion of radial glia into basal precursors and guides neuronal amplification by indirect neurogenesis in the developing neocortex. *Neuron*, *60*(1), 56–69. <https://doi.org/10.1016/j.neuron.2008.09.028>
- Shen, J., Lu, J., Sui, L., Wang, D., Yin, M., Hoffmann, I., ... Pflugfelder, G. O. (2014). The orthologous Tbx transcription factors Omb and TBX2 induce epithelial cell migration and extrusion in vivo without involvement of matrix metalloproteinases. *Oncotarget*, *5*(23), 11998–12015. <https://doi.org/10.18632/oncotarget.2426>
- Sonoda, T., Lee, S. K., Birnbaumer, L., & Schmidt, T. M. (2018). Melanopsin phototransduction is repurposed by ipRGC subtypes to shape the function of distinct visual circuits. *Neuron*, *99*(4), 754–767 e754. <https://doi.org/10.1016/j.neuron.2018.06.032>
- Sonoda, T., Okabe, Y., & Schmidt, T. M. (2019). Overlapping morphological and functional properties between M4 and M5 intrinsically photosensitive retinal ganglion cells. *The Journal of Comparative Neurology*, *528*, 1028–1040. <https://doi.org/10.1002/cne.24806>
- Stabio, M. E., Sabbah, S., Quattrocchi, L. E., Ilardi, M. C., Fogerson, P. M., Leyrer, M. L., Kim, M. T., Kim, I., Schiel, M., Renna, J. M., Briggman, K. L., & Berson, D. M. (2018). The M5 cell: A color-opponent intrinsically photosensitive retinal ganglion cell. *Neuron*, *97*(1), 251. <https://doi.org/10.1016/j.neuron.2017.12.030>
- Strumpf, D., Mao, C. A., Yamanaka, Y., Ralston, A., Chawengsaksophak, K., Beck, F., & Rossant, J. (2005). Cdx2 is required for correct cell fate specification and differentiation of trophectoderm in the mouse blastocyst. *Development*, *132*(9), 2093–2102. <https://doi.org/10.1242/dev.01801>
- Sweeney, N. T., Tierney, H., & Feldheim, D. A. (2014). Tbr2 is required to generate a neural circuit mediating the pupillary light reflex. *The Journal of Neuroscience*, *34*(16), 5447–5453. <https://doi.org/10.1523/JNEUROSCI.0035-14.2014>
- Tran, N. M., Shekhar, K., Whitney, I. E., Jacobi, A., Benhar, I., Hong, G., Yan, W., Adiconis, X., Arnold, M. K. E., Lee, J. M., Levin, J. Z., Lin, D., Wang, C., Lieber, C. M., Regev, A., He, Z., & Sanes, J. R. (2019). Single-cell profiles of retinal ganglion cells differing in resilience to injury reveal neuroprotective genes. *Neuron*, *104*(6), 1039–1055 e1012. <https://doi.org/10.1016/j.neuron.2019.11.006>
- Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., Jr., & Lowell, B. B. (2011). Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron*, *71*(1), 142–154. <https://doi.org/10.1016/j.neuron.2011.05.028>
- Wu, F., Kaczynski, T. J., Sethuramanujam, S., Li, R., Jain, V., Slaughter, M., & Mu, X. (2015). Two transcription factors, Pou4f2 and Isl1, are sufficient to specify the retinal ganglion cell fate. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(13), E1559–E1568. <https://doi.org/10.1073/pnas.1421535112>
- Xiang, M., Zhou, L., Macke, J. P., Yoshioka, T., Hendry, S. H., Eddy, R. L., Shows, T. B., & Nathans, J. (1995). The Brn-3 family of POU-domain factors: Primary structure, binding specificity, and expression in subsets of retinal ganglion cells and somatosensory neurons. *The Journal of Neuroscience*, *15*(7 Pt 1), 4762–4785.
- Xue, T., Do, M. T., Riccio, A., Jiang, Z., Hsieh, J., Wang, H. C., ... Yau, K. W. (2011). Melanopsin signalling in mammalian iris and retina. *Nature*, *479*(7371), 67–73. <https://doi.org/10.1038/nature10567>
- Yan, W., Laboulaye, M. A., Tran, N. M., Whitney, I. E., Benhar, I., & Sanes, J. R. (2020). Mouse retinal cell atlas: Molecular identification of over sixty amacrine cell types. *J Neurosci.*, *40*(27), 5177–5195. <https://doi.org/10.1523/JNEUROSCI.0471-20.2020>

How to cite this article: Chen, C.-K., Kiyama, T., Weber, N., Whitaker, C. M., Pan, P., Badea, T. C., Massey, S. C., & Mao, C.-A. (2021). Characterization of Tbr2-expressing retinal ganglion cells. *Journal of Comparative Neurology*, *529*(15), 3513–3532. <https://doi.org/10.1002/cne.25208>